TABLE 3.—DISTRIBUTION OF POLLOCK IN THE CENTRAL AND WESTERN REGULATORY AREAS OF THE GULF OF ALASKA. SEASONAL BIOMASS DISTRIBUTION, AREA APPORTIONMENTS, AND SEASONAL ALLOWANCES OF ANNUAL TAC IN 2004

[Values are in metric tons]						
Season	Shumagin (Area 610) (biomass distribution)	Chirikof (Area 620) (biomass distribution)	Kodiak (Area 630) (biomass distribution)	Total (biomass distribution)		
A	3,747 (23.63%)	9,027 (56.9%)	3,091 (19.48%)	15,865 (100%)		

Season	(biomass distribution)	(biomass distribution)	(biomass distribution)	(biomass distribution)
A	3,747 (23.63%)	9,027 (56.9%)	3,091 (19.48%)	15,865 (100%)
В	3,748 (23.62%)	10,704 (67.47%)	1,413 (8.91%)	15,865 (100%)
C	7,717 (48.64%)	3,380 (21.3%)	4,768 (30.06%)	15,865 (100%)
D	7,718 (48,64%)	3,379 (21.3%)	4,768 (30.06%)	15,865 (100%)

22.930

26,490

14.040

63,460

Season	(Area 610) (biomass distribution)	(Area 620) (biomass distribution)	(Area 630) (biomass distribution)	Total (biomass distribution)
A	3,747 (23.63%)	9,027 (56.9%)	3,091 (19.48%)	15,865 (100%)
В	3,748 (23.62%)	10,704 (67.47%)	1,413 (8.91%)	15,865 (100%)
C	7,717 (48.64%)	3,380 (21.3%)	4,768 (30.06%)	15,865 (100%)